



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

EP04/111 04

REC'D 01 DEC 2004

WIPO

PCT

Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

03023342.3

**PRIORITY
DOCUMENT**
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk

BEST AVAILABLE COPY



Anmeldung Nr:
Application no.: 03023342.3
Demande no:

Anmeldetag:
Date of filing: 15.10.03
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Newron Pharmaceuticals S.p.A.
Via L. Ariosto, 21
20091 Bresso (MI)
ITALIE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Substituted benzylaminoalkylene heterocycles

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

C07D307/00

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL
PT RO SE SI SK TR LI

sodium and/or calcium channel modulators and/or as selective MAO-B inhibitors and therefore useful in preventing, alleviating and curing a wide range of pathologies, including, but not limited to, neurological, psychiatric, cardiovascular, inflammatory, ophthalmic, urologic, metabolic and
5 gastrointestinal diseases, where the above mechanisms have been described as playing a pathological role.

BACKGROUND OF THE INVENTION

Chemical background

WO 02/49993, discloses aryl-substituted heterocyclic derivatives and
10 the use of such compounds in treating a variety of inflammatory and immuno system disorders. US patent 4,145,439 describes benzyl phenyl ethers and their use for combatting arthropodes, nematodes, fungi and bacteria. WO 03/37865 describes the preparation of pyridinylethyl amines and amides and their use as anticancer drugs. WO 01/14331 describes the preparation of
15 4-amino-1-benzylpiperidines(piperazines) as antimalarial agents. In US patent 5,318,988 2-aminomethyl-chromanes are described together with their use in the treatment of CNS diseases. In J. Indian Chem. Soc. 1987, 64, 169-171, the synthesis of benzodioxolano-ethylamines is reported. In Armyanskii Khimicheskii Zhurnal 1968, 21, 509-514, the synthesis of benzodioxan
20 derivatives is described as possible adrenolytic substances. In WO 96/20191, the preparation of 3-(N-aryl and N-heterocyclylaminomethyl)-indole derivatives are described and their use against CNS diseases. In FR patent 2.181.559 indole derivatives are described and their use as anticonvulsant and analgesic agents. In WO 94/26738 benzofuranylurea derivatives and analogues
25 were described as ACAT-inhibitors. In EP 534246 benzylaminoderivatives are described and their use as antiarrhythmic agents. In WO 01/12604, 2-pyridyl derivatives are described as fungicides.

Biological background

It is well known that sodium channels play an important role in the neuronal network by transmitting electrical impulses rapidly throughout cells and cell networks, thereby coordinating higher processes ranging from locomotion to cognition. These channels are large transmembrane proteins, which are able to switch between different states to enable selective permeability for sodium ions. For this process an action potential is needed to depolarize the membrane, and hence these channels are voltage-gated. In the past few years a much better understanding of sodium channels and drugs interacting with them has been developed.

It has become clear that a number of drugs having an unknown mechanism of action actually act by modulating sodium channel conductance, including local anesthetics, class I antiarrhythmics and anticonvulsants. Neuronal sodium channel blockers have found application with their use in the treatment of epilepsy (phenytoin and carbamazepine), bipolar disorder (lamotrigine), preventing neurodegeneration, and in reducing neuropathic pain. Various anti-epileptic drugs that stabilize neuronal excitability are effective in neuropathic pain (gabapentin).

In addition, an increase in sodium channel expression or activity has also been observed in several models of inflammatory pain, suggesting a role of sodium channels in inflammatory pain.

Calcium channels are membrane-spanning, multi-subunit proteins that allow controlled entry of calcium ions into cells from the extracellular fluid. Commonly, calcium channels are voltage dependent and are referred to as voltage sensitive calcium channels (VSCC). VSCCs are found throughout the mammalian nervous system, where they regulate such varied activities as cellular excitability, transmitter release, intracellular metabolism, neurosecretory activity and gene expression. All "excitable" cells in animals,

such as neurons of the central nervous system (CNS), peripheral nerve cells, and muscle cells, including those of skeletal muscles, cardiac muscles and venous and arterial smooth muscles, have voltage dependent calcium channels. Calcium channels have a central role in regulating intracellular calcium ions levels that are important for cell viability and function. Intracellular calcium ion concentrations are implicated in a number of vital processes in animals, such as neurotransmitter release, muscle contraction, pacemaker activity, and secretion of hormones. It is believed that calcium channels are relevant in certain disease states. A number of compounds useful in treating various cardiovascular diseases in mammals, including humans, are thought to exert their beneficial effects by modulating functions of voltage dependant calcium channels present in cardiac and/or vascular smooth muscle. Compounds with activity against calcium channels have also been implicated for the treatment of pain. In particular N-type calcium channels (Cav2.2), responsible for the regulation of neurotransmitters, are thought to play a significant role in nociceptive transmission, both due to their tissue distribution as well as from the results of several pharmacological studies. This hypothesis has been validated in the clinic by Zinocotide, a peptide derived from the venom of the marine snail, *Conus Magus*. A limitation in the therapeutic use of this peptide is that it has to be administered intrathecally in humans (Bowersox S. S. and Luther R. *Toxicon*, 1998, 36, 11, 1651-1658).

Monoamine oxidase (MAO) is an integral protein of the outer mitochondrial membrane and plays a major role in the in vivo inactivation of biogenic and diet-derived amines in both the CNS and in peripheral neurons and tissues. Two MAO enzymes are distinguished on the basis of their substrate preferences and sensitivity to inhibition by the MAO inhibitor, clorgyline:

- MAO type A (MAO-A) which in the human CNS is responsible for the deamination of serotonin and noradrenaline. The highest MAO-A concentrations are in the catecholaminergic neurons of the locus ceruleus;

- MAO-B which is responsible mainly for the catabolism of dopamine.

5 In contrast to the rat brain, MAO-B is the major form in the human and guinea pig CNS. The highest MAO-B concentrations are in the serotonergic neurons of the raphe nucleus and posterior hypothalamus. The nigral MAO-B is located primarily in glial cells.

MAO-B (but not MAO-A) activity in CNS increases with age in both
10 humans and animals, perhaps as a result of the glial cell proliferation associated with neuronal loss. Increased MAO-B levels in Alzheimer's plaques have also been reported. Increased blood platelet MAO-B activity has been reported in both Alzheimer's (AD) and Parkinson's Disease (PD). MAO-B activity was reduced by 40% in the brain of smokers: tobacco use is
15 associated both with psychiatric disease and with a reduced risk for PD.

Most currently investigated MAO-B inhibitors are irreversible. The inhibition is very persistent (weeks), as its effects can only be overcome by de novo synthesis of the enzyme. Interest in the MAO-B inhibition was initially stimulated by the desire to elevate the reduced striatal DA concentrations
20 characteristic of PD. As increased DA concentration in the synaptic cleft would be expected as the primary effect of treatment with a selective MAO-B inhibitor. In PD, the need to supply the DA precursor L-Dopa, the golden standard PD therapy, should thus be reduced. This is desirable, as L-Dopa, despite the excellent initial improvement achieved, is associated in the longer
25 term with increasingly severe side effects, including motor fluctuations, dyskinesia and dystonia.

In addition to loss of cholinergic neurons, there is a decrease in the levels of the DA, noradrenaline and serotonin in the brains of AD patients.

MAO-B activity is significantly increased in the platelets and selected brain regions of AD patients. MAO-B inhibitors may act by both reducing the formation of oxygen radicals and preventing the breakdown of monoamines and thus elevating their level in the brain of AD patients.

5 Promising results using MAO-B inhibitors were reported also for the treatment of narcolepsy, Tourette's syndrome/attention deficit disorders, negative symptoms of schizophrenia, drug addiction and obesity.

All together these findings indicate that compounds with sodium and/or calcium channel blockade and/or MAO-B inhibition have a high therapeutic
10 potential in preventing, alleviating and curing a wide range of pathologies, including neurological, psychiatric, cardiovascular, urologic, metabolic and gastrointestinal diseases, where the above mechanisms have been described as playing a pathological role.

There are many papers and patents which describe sodium channel
15 and/or calcium channel modulators or antagonists and/or MAO-B inhibitors for the treatment or modulation of a plethora of disorders, such as their use as local anesthetics, antiarrhythmics, antiemetics, antimanic depressants, agents for the treatment of unipolar depression, cardiovascular diseases, urinary incontinence, diarrhea, inflammation, epilepsy, neurodegenerative conditions,
20 nerve cell death, anticonvulsants, neuropathic pain, migraine, acute hyperalgesia and inflammation, renal disease, allergy, asthma, bronchospasm, dysmenorrhea, esophageal spasm, glaucoma, urinary tract disorders, gastrointestinal motility disorders, premature labour, obesity. A largely incomplete list is shown below.


25 An extensive and thorough prior art overview is reported in WO 03/057219 (and references therein); a further selection of prior art is reported in the following references: Alzheimer, C. Adv. Exp. Med. Biol. 2002, 513, 161-181; Vanegas, H.; Schaible, H. Pain 2000, 85, 9-18; U.S. Patent

7

5,051,403; U.S. Patent 5,587,454; U.S. Patent 5,863,952; U.S. Patent
 6,011,035; U.S. Patent 6,117,841; U.S. Patent 6,362,174; U.S. Patent
 6,380,198; U.S. Patent 6,420,383; U.S. Patent 6,458,781; U.S. Patent
 6,472,530; U.S. Patent 6,518,288; U.S. Patent 6,521,647; WO 97/10210; WO
 5 03/018561; WO 01/12176; WO 96/37199; U.S. patent 6, 210,706.

DESCRIPTION OF THE INVENTION

This invention is related to benzylaminomethylene heterocycles of the following general formula (I)



I

wherein

X is oxygen or sulphur or a NR⁷ group;

R¹ is C₃-C₈ alkyl, or C₁-C₈ alkyl substituted by phenoxy or phenyl,
 both phenoxy or phenyl being optionally substituted by one or
 more fluoro, chloro, trifluoromethyl, C₁-C₆ alkyl, hydroxyl,
 C₁-C₆ alkoxy;

R², R³ are independently hydrogen, C₁-C₆ alkyl, halogen,
 trifluoromethyl, hydroxy or C₁-C₆ alkoxy;

R⁴ is hydrogen, C₁-C₈ alkyl;

R⁵, R⁶ are independently hydrogen, C₁-C₃ alkyl, optionally substituted
 by hydroxy or phenyl;

R⁷ is hydrogen or straight or branched C₁-C₃ alkyl;

Het is a five to seven membered, saturated or unsaturated
 heteromonocyclic or an eight to ten membered, saturated or

unsaturated heterobicyclic group, containing one or more heteroatoms chosen independently from nitrogen, oxygen and sulphur, said mono- or bicyclic groups being optionally substituted by C₁-C₆alkyl, halogen, hydroxyl or C₁-C₆ alkoxy;

5 and the pharmaceutically acceptable salts thereof, with the proviso that:


- Het cannot be indole (FR 2,181,559), benzo[b]furan (WO 94/26738), benzo[b]thiophen (WO 94/26738), chroman (US 5,318,988), when R⁵ and R⁶ are both hydrogen, or 2-pyridyl (WO 01/12604);
- 10 • when R¹ is unsubstituted or substituted benzyl, and R² and R³ are hydrogen, halogen or alkoxy, R⁴ is other than hydrogen (US 4,145,439);
- when R¹ is propyl or butyl, R², R³, R⁵ and R⁶ are hydrogen and R⁴ is hydrogen, methyl or ethyl, Het cannot be 1,4-benzodioxan (Mndzhoyan *et al.* Armyanskii Khimicheskii Zhurnal 1968, 21, 509-514);
- 15 • when X- R¹ is a para butyloxy group and R², R³, R⁴, R⁵ and R⁶ are hydrogen, Het cannot be 2-thiophenyl, (US 4,623,662) or 4-(2,2'-dimethyl)-pyranyl (Arutyunian, N. S. et al. Armyanskii Khimicheskii Zhurnal 1984, 37, 749-753);
- 20 • when X- R¹ is an ortho heptyloxy or octyloxy group and R², R³, R⁴, R⁵ and R⁶ are hydrogen, Het cannot be 2-furyl (commercial library)
- when X- R¹ is an ortho O-(CH₂)_m-p-CF₃-phenyl group, where m is an integer from 1 to 3, and R², R³, R⁵ and R⁶ are hydrogen, Het cannot be pyridyl (EP 534246)
- 25

The optional substituents in the phenyl ring may be in any position. The pharmaceutically acceptable salts of the compounds of formula I include acid

9

addition salts with inorganic acids, e.g. hydrochloric, hydrobromic, sulphuric, and phosphoric or organic acids, e.g. acetic, propionic, benzoic, cinnamic, mandelic, salicylic, glycolic, lactic, oxalic, malic, maleic, malonic, fumaric, tartaric, citric, and the like.

5 The invention also relates to the use of the compounds of formula I



I

wherein

- 10 X is oxygen or sulphur or a NR⁷ group;
- R¹ is C₃-C₈ alkyl, or C₁-C₈ alkyl substituted by phenoxy or phenyl, both phenoxy or phenyl being optionally substituted by one or more fluoro, chloro, trifluoromethyl, C₁-C₆ alkyl, hydroxyl, C₁-C₆ alkoxy;
- 15 R², R³ are independently hydrogen, C₁-C₆ alkyl, halogen, trifluoromethyl, hydroxy or C₁-C₆ alkoxy;
- R⁴ is hydrogen, C₁-C₈ alkyl;
- R⁵, R⁶ are independently hydrogen, C₁-C₃ alkyl, optionally substituted by hydroxy or phenyl;
- 20 R⁷ is hydrogen or straight or branched C₁-C₃ alkyl;
- Het is a five to seven membered, saturated or unsaturated heteromonocyclic or an eight to ten membered, saturated or unsaturated heterobicyclic group, containing one or more heteroatoms chosen independently from nitrogen, oxygen and
- 25 sulphur, said mono- or bicyclic groups being optionally

substituted by C₁-C₆alkyl, halogen, hydroxyl or C₁-C₆ alkoxy;
and the pharmaceutically acceptable salts thereof, with the proviso that:

- Het cannot be an indole (FR 2,181,559), chroman (US 5,318,988), when R⁵ and R⁶ are both hydrogen;
- 5 • when X- R¹ is an ortho O-(CH₂)_m-p-CF₃-phenyl group, where m is an integer from 1 to 3, and R², R³, R⁵ and R⁶ are hydrogen, Het cannot be pyridyl (EP 534246)

for the preparation of a medicament having sodium and/or calcium channel modulating activity and/or selective MAO-B inhibiting activity.

- 10 Preferred use compounds of the invention are the compounds of formula I wherein R¹ is benzyl or C₅-C₈ alkyl, R⁴, R⁵, and R⁶ are hydrogen or C₁-C₃ alkyl, X is oxygen and Het is furan, tetrahydrofuran, isoxazol, oxazol, thiophen, pyran, dioxane, unsubstituted or substituted by C₁-C₃ alkyl.

Examples of specific compounds of the invention are:

- 15 (4-Pentyloxy-benzyl)-(furan-2-ylmethyl)-amine;
 (4-Heptyloxy-benzyl)-(furan-2-ylmethyl)-amine;
 (R) (4-Pentyloxy-benzyl)-(tetrahydro-furan-2-ylmethyl)-amine;
 (S) (4-Pentyloxy-benzyl)-(tetrahydro-furan-2-ylmethyl)-amine;
 (R) (4-Heptyloxy-benzyl)-(tetrahydro-furan-2-ylmethyl)-amine;
20 (S) (4-Heptyloxy-benzyl)-(tetrahydro-furan-2-ylmethyl)-amine;
 (R) (4-Pentyloxy-benzyl)-(tetrahydro-pyran-2-ylmethyl)-amine;
 (S) (4-Pentyloxy-benzyl)-(tetrahydro-pyran-2-ylmethyl)-amine;
 (R) (4-Heptyloxy-benzyl)-(tetrahydro-pyran-2-ylmethyl)-amine;
 (S) (4-Heptyloxy-benzyl)-(tetrahydro-pyran-2-ylmethyl)-amine;
25 (4-Benzyloxy-benzyl)-(furan-2-ylmethyl)-amine;
 (4-Benzyloxy-benzyl)-(5-methyl-furan-2-ylmethyl)-amine;
 [4-(3-Fluoro-benzyloxy)-benzyl]-(furan-2-ylmethyl)-amine;
 [4-(3-Chloro-benzyloxy)-benzyl]-(furan-2-ylmethyl)-amine;


- [4-(3-Fluoro-benzyloxy)-benzyl]-(5-methyl-furan-2-ylmethyl)-amine;
[4-(3-Chloro-benzyloxy)-benzyl]-(5-methyl-furan-2-ylmethyl)-amine;
(R) (4-Benzyloxy-benzyl)-[1-(furan-2-yl)-1-ethyl]-amine;
(S) (4-Benzyloxy-benzyl)-[1-(furan-2-yl)-1-ethyl]-amine;
5 [4-(3-Fluoro-benzyloxy)-benzyl]-[1-(furan-2-yl)-1-ethyl]-amine;
[4-(3-Chloro-benzyloxy)-benzyl]-[1-(furan-2-yl)-1-ethyl]-amine;
(R) (4-Benzyloxy-benzyl)-(tetrahydro-furan-2-ylmethyl)-amine;
(S) (4-Benzyloxy-benzyl)-(tetrahydro-furan-2-ylmethyl)-amine;
(R) [4-(3-Fluoro-benzyloxy)-benzyl]-(tetrahydro-furan-2-ylmethyl)-amine;
10 (S) [4-(3-Fluoro-benzyloxy)-benzyl]-(tetrahydro-furan-2-ylmethyl)-amine;
(R) [4-(3-Chloro-benzyloxy)-benzyl]-(tetrahydro-furan-2-ylmethyl)-amine;
(S) [4-(3-Chloro-benzyloxy)-benzyl]-(tetrahydro-furan-2-ylmethyl)-amine;
(R) [4-(3-Fluoro-benzyloxy)-benzyl]-(tetrahydro-pyran-2-ylmethyl)-amine;
(S) [4-(3-Fluoro-benzyloxy)-benzyl]-(tetrahydro-pyran-2-ylmethyl)-amine;
15 (R) [4-(3-Chloro-benzyloxy)-benzyl]-(tetrahydro-pyran-2-ylmethyl)-amine;
(S) [4-(3-Chloro-benzyloxy)-benzyl]-(tetrahydro-pyran-2-ylmethyl)-amine;
(4-Benzyloxy-benzyl)-(1,4-dioxan-2-ylmethyl)-amine;
[4-(3-Chloro-benzyloxy)-benzyl]-(pyrido-2-ylmethyl)-amine;
[4-(3-Chloro-benzyloxy)-benzyl]-(pyrido-3-ylmethyl)-amine;
20 [4-(3-Chloro-benzyloxy)-benzyl]-(pyrido-4-ylmethyl)-amine;
[4-(3-Chloro-benzyloxy)-benzyl]-(imidazol-2-ylmethyl)-amine;
[4-(3-Chloro-benzyloxy)-benzyl]-(1-methyl-imidazol-2-ylmethyl)-amine;
[4-(3-Chloro-benzyloxy)-benzyl]-(5-methyl-1H-triazol-2-ylmethyl)-amine;
[4-(3-Chloro-benzyloxy)-benzyl]-(4-methyl-thiazol-2-ylmethyl)-amine;
25 (4-Benzyloxy-benzyl)-(isoxazol-5-ylmethyl)-amine;
[4-(3-Fluoro-benzyloxy)-benzyl]-(isoxazol-5-ylmethyl)-amine;
[4-(3-Chloro-benzyloxy)-benzyl]-(isoxazol-5-ylmethyl)-amine;
(4-Benzyloxy-benzyl)-(3-methyl-isoxazol-5-ylmethyl)-amine;

- [4-(3-Fluoro-benzyloxy)-benzyl]-(3-methyl-isoxazol-5-ylmethyl)-amine;
[4-(3-Chloro-benzyloxy)-benzyl]-(3-methyl-isoxazol-5-ylmethyl)-amine;
(4-Benzyloxy-benzyl)-(oxazol-2-ylmethyl)-amine;
[4-(3-Fluoro-benzyloxy)-benzyl]-(oxazol-2-ylmethyl)-amine;
5 [4-(3-Chloro-benzyloxy)-benzyl]-(oxazol-2-ylmethyl)-amine;
(4-Benzyloxy-benzyl)-(oxazol-5-ylmethyl)-amine;
[4-(3-Fluoro-benzyloxy)-benzyl]-(oxazol-5-ylmethyl)-amine;
[4-(3-Chloro-benzyloxy)-benzyl]-(oxazol-5-ylmethyl)-amine;
(4-Benzyloxy-benzyl)-(thiophen-2-ylmethyl)-amine;
10 [2-(3-Chloro-benzyloxy)-benzyl]-(thiophen-2-ylmethyl)-amine;
[2-(3-Fluoro-benzyloxy)-benzyl]-(thiophen-2-ylmethyl)-amine;
[3-(3-Chloro-benzyloxy)-benzyl]-(thiophen-2-ylmethyl)-amine;
[3-(3-Fluoro-benzyloxy)-benzyl]-(thiophen-2-ylmethyl)-amine;
[4-(3-Chloro-benzyloxy)-benzyl]-(thiophen-2-ylmethyl)-amine;
15 [4-(3-Fluoro-benzyloxy)-benzyl]-(thiophen-2-ylmethyl)-amine;
[2-(3-Fluoro-benzyloxy)-benzyl]-(benzo[b]furan-2-ylmethyl)-amine;
[3-(3-Fluoro-benzyloxy)-benzyl]-(benzo[b]furan-2-ylmethyl)-amine;
[4-(3-Fluoro-benzyloxy)-benzyl]-(benzo[b]furan-2-ylmethyl)-amine;
(R) [4-(3-Fluoro-benzyloxy)-benzyl]-(dihydro-benzo[b]furan-2-ylmethyl)-
20 amine;
(S) [4-(3-Fluoro-benzyloxy)-benzyl]-(dihydro-benzo[b]furan-2-ylmethyl)-
amine;
[4-(3-Chloro-benzyloxy)-benzyl]-(benzimidazol-2-ylmethyl)-amine;
either as a single isomer or as a mixture of isomers thereof, and the
25 pharmaceutically acceptable salts thereof.

The compounds of the invention and the salts thereof can be obtained
by a process comprising:


13

a) reaction of compounds of formula II



II


5 wherein R^1 , R^2 , R^3 and X are as defined above
with compounds of formula III, in the presence of a reducing agent



III

10 wherein R^4 , R^5 , R^6 and Het are as defined previously thus obtaining a
compound of formula I; or

b) reaction of compounds of formula IV



IV

15

wherein R^1 , R^2 , R^3 and X are as defined above and Y is a halogen atom
or a O-EWG group, where the EWG means an electron withdrawing group,
like e.g. mesyl, tosyl or trifluoroacetyl groups, able to transform the oxygen
which they are linked to, in a good leaving group with compounds of formula
20 III and, if desired, converting a compound of the invention into another
compound of the invention and/or, if desired, converting a compound of the
invention into a pharmaceutically acceptable salt and/or, if desired, converting

a salt into a free compound.

Compounds II-IV are commercially available compounds or are prepared from commercially available compounds using well-known methods.

The reaction compounds of formula II with compounds of formula III to give the compounds of formula I is a reductive amination reaction which can be carried out according to well known methods. According to a preferred embodiment of the invention, it may be performed under nitrogen atmosphere, in a suitable organic solvent, such as an alcohol, e.g. a lower alkanol, in particular methanol, or in acetonitrile, or in tetrahydrofuran, at a temperature ranging from about 0°C to about 70°C, in the presence of a reducing agent, the most appropriate being sodium borohydride or sodium cyanoborohydride. Occasionally titanium IV isopropylate and molecular sieves can be added to the reaction mixture for facilitating the reaction.

In a compound of formula IV the halogen is preferably bromine or iodine. The alkylation reaction of a compound of formula IV with a compound of formula III can be carried out in a suitable organic solvent, such as an alcohol, e.g. methanol, ethanol or isopropanol, in particular in ethanol, at a temperature ranging from about 0°C to about 50°C.

The optional salification of a compound of the invention as well as the conversion of a salt into the free compound may be carried out by conventional methods.

When in the compounds of the present invention and in the intermediate-products thereof, groups are present, which need to be protected before submitting them to the above illustrated reactions, they may be protected before being reacted and then deprotected according to well-known methods.

PHARMACOLOGY

The compounds of the invention display affinities for the calcium

15

and/or sodium channel binding sites as demonstrated using selective radioligands in *in vitro* binding studies.

The compounds according to the invention are blockers of the voltage dependent sodium channels and/or calcium channels. These compounds
5 therefore displace 3H-batrachotoxin (BTX) with a high affinity from the binding site on the sodium channel, with IC_{50} in the low μM or sub μM range. Similarly the compounds displace 3H-nitrendipine from the binding site in the calcium channel, with IC_{50} in the low μM or most usually in the sub μM range as well as inhibiting calcium influx induced through calcium channels via
10 cellular depolarisation.

Such substances exhibit "use-dependency" when the sodium channels are blocked i.e. maximum blockage of the sodium channels is only achieved after repeated stimulation of the sodium channel. Consequently, the substances preferably bind to sodium channels which are multiply activated.
15 As a result the substances are capable of activity preferentially in those regions of the body which are pathologically over-stimulated, as illustrated by patch-clamp experiments (W.A. Catterall, Trends Pharmacol. Sci., 8, 57-65; 1987) which show that the compounds according to the invention block the electrically stimulated sodium channel in a "use-dependent" manner.

20 As a consequence of these mechanisms the compounds of the invention are active *in vivo* when orally administered in the range of 0.1 to 100 mg/kg in a wide range of animal models and in particular in the MES test of electrically-induced convulsions, in the formalin model of persistent pain and in the carrageenan model of inflammation.

25 The compounds of this invention beyond being potent calcium and/or sodium channels blockers are also able to selectively and reversibly inhibit MAO-B both *in vitro* and *in vivo*.

The compounds of the invention that are potent inhibitors of MAO-B

(IC₅₀ in the submicromolar range) have generally no relevant effect on MAO-A. In enzymatic studies using rat brain membranes and selective radiolabelled substrates, the compounds are generally over 1000 times more active as MAO-B inhibitors than MAO-A inhibitors. The MAO-B inhibition
5 by the compounds is not time-dependent and the enzyme activity fully recovers after washings as it is typical for reversible inhibitors.

After oral administration in mice, the compounds behave as potent and reversible, short-acting MAO-B inhibitors with full recovery of activity 24 h after treatment.

10 In view of the above described mechanisms of action, the compounds of the present invention are useful in the treatment or prevention of neuropathic pain. Neuropathic pain syndromes include, and are not limited to: diabetic neuropathy; sciatica; non-specific lower back pain; multiple sclerosis pain; fibromyalgia; HIV-related neuropathy; neuralgia, such as post-herpetic
15 neuralgia and trigeminal neuralgia; and pain resulting from physical trauma, amputation, cancer, toxins or chronic inflammatory conditions.

Compounds of the invention are also useful for the treatment of chronic pain. Chronic pain includes, and is not limited to, chronic pain caused by inflammation or an inflammatory-related condition, osteoarthritis,
20 rheumatoid arthritis or as sequela to disease, acute injury or trauma and includes upper back pain or lower back pain (resulting from systematic, regional or primary spine disease (such as radiculopathy), bone pain (due to osteoarthritis, osteoporosis, bone metastasis or unknown reasons), pelvic pain, spinal cord injury-associated pain, cardiac chest pain, non-cardiac chest pain,
25 central post-stroke pain, myofascial pain, cancer pain, AIDS pain, sickle cell pain, geriatric pain or pain caused by headache, temporomandibular joint syndrome, gout, fibrosis or thoracic outlet syndromes.

Compounds of the invention are also useful in the treatment of acute

pain (caused by acute injury, illness, sports-medicine injuries, carpal tunnel syndrome, burns, musculoskeletal sprains and strains, musculotendinous strain, cervicobrachial pain syndromes, dyspepsia, gastric ulcer, duodenal ulcer, dysmenorrhea, endometriosis or surgery (such as open heart or bypass surgery), post operative pain, kidney stone pain, gallbladder pain, gallstone pain, obstetric pain or dental pain.

Compounds of the invention are also useful in the treatment of migraine, such as tension type headache, transformed migraine or evolutive headache, cluster headache, as well as secondary headache disorders, such as the ones derived from infections, metabolic disorders or other systemic illnesses and other acute headaches, paroxysmal hemicrania and the like, resulting from a worsening of the above mentioned primary and secondary headaches.

Compounds of the invention are also useful for the treatment of neurological conditions and cognitive disorders. Neurological conditions include, and are not limited to, conditions such as epilepsy (including simple partial seizures, complex partial seizures, secondary generalised seizures, further including absence seizures, myoclonic seizures, clonic seizures, tonic seizures, tonic clonic seizures and atonic seizures), degenerative dementia (including senile dementia, Alzheimer's disease, Pick's disease, Huntington's chorea, Parkinson's disease and Creutzfeldt-Jakob disease), and vascular dementia (including multi-infarct dementia, stroke and cerebral ischemia), as well as dementia associated with intracranial space occupying lesions, trauma, infections and related conditions (including HIV infection), metabolism, toxins, anoxia and vitamin deficiency; and mild cognitive impairment associated with aging, particularly age associated memory impairment, movement disorders (postencephalitic parkinsonism, progressive supranuclear palsy, corticobasal degeneration, restless leg syndrome), narcolepsy, deficit and hyperactivity

disorders (ADHD), Tourette's Syndrome, amyotrophic lateral sclerosis, Down's syndrome.

Compounds of the invention are also useful for the treatment of psychiatric disorders. Psychiatric disorders include, and are not limited to, manic depression also known as bipolar disorder (such as bipolar disorder type I, bipolar disorder type II), cyclothymic disorder, rapid cycling, ultradian cycling, bipolar depression, acute mania, mania, mixed mania, hypomania or unipolar depression, schizophrenia, schizophreniform disorders, schizoaffective disorders, delusional disorders, brief psychotic disorders, shared psychotic disorders, psychotic disorder due to a general medical condition, substance-induced psychotic disorders or a psychotic disorder not otherwise specified, anxiety disorders and moreover in smoke and drug addiction.

Compounds of the invention are also useful in the treatment of peripheral diseases such as tinnitus, muscle spasm, muscular sclerosis, and other disorders, including, and not limited to cardiovascular diseases (such as cardiac arrhythmia, cardiac infarction or angina pectoris, hypertension, hypoxia, cardiac ischemia) endocrine disorders (such as acromegaly or diabetes insipidus), diseases in which the pathophysiology of the disorder involves excessive or hypersecretory or otherwise inappropriate cellular secretion of an endogenous substance (such as catecholamine, a hormone or a growth factor).

Compounds of the invention are also useful in the treatment of liver disease, such as inflammatory liver disease, for example chronic viral hepatitis B, chronic viral hepatitis C, alcoholic liver injury, primary biliary cirrhosis, autoimmune hepatitis, non-alcoholic steatohepatitis and liver transplant rejection.

Compounds of the invention inhibit inflammatory processes affecting

all body systems. Therefore are useful in the treatment of inflammatory processes of the muscular-skeletal system of which the following is a list of examples but it is not comprehensive of all target disorders: arthritic conditions such as ankylosing spondylitis, cervical arthritis, fibromyalgia, gut, 5 juvenile rheumatoid arthritis, lumbosacral arthritis, osteoarthritis, osteoporosis, psoriatic arthritis, rheumatic disease; disorders affecting skin and related tissues: eczema, psoriasis, dermatitis and inflammatory conditions such as sunburn; disorders of the respiratory system: asthma, allergic rhinitis and respiratory distress syndrome, lung disorders in which inflammation is 10 involved such as asthma and bronchitis; chronic obstructive pulmonary disease; disorders of the immune and endocrinological systems: periarteritis nodosa, thyroiditis, aplastic anaemia, scleroderma, myasthenia gravis, multiple sclerosis, sarcoidosis, nephritic syndrome, Bechet's syndrome, polymyositis, gingivitis.

15 Compounds of the invention are also useful in the treatment of gastrointestinal (GI) tract disorders such as inflammatory bowel disorders including but not limited to ulcerative colitis, Crohn's disease, ileitis, proctitis, celiac disease, enteropathies, microscopic or collagenous colitis, eosinophilic gastroenteritis, or pouchitis resulting after proctocolectomy and 20 post ileonatal anastomosis, and irritable bowel syndrome including any disorders associated with abdominal pain and/or abdominal discomfort such as pylorospasm, nervous indigestion, spastic colon, spastic colitis, spastic bowel, intestinal neurosis, functional colitis, mucous colitis, laxative colitis and functional dyspepsia; but also for treatment of atrophic gastritis, gastritis 25 varioliforme, ulcerative colitis, peptic ulceration, pyresis, and other damage to the GI tract, for example, by *Helicobacter pylori*, gastroesophageal reflux disease, gastroparesis, such as diabetic gastroparesis; and other functional bowel disorders, such as non-ulcerative dyspepsia (NUD); emesis, diarrhea,

and visceral inflammation.

Compounds of the invention are also useful in the treatment of disorders of the genito-urinary tract such as overactive bladder, prostatitis (chronic bacterial and chronic non-bacterial prostatitis), prostatic dysnia, interstitial
5 cystitis, urinary incontinence and benign prostatic hyperplasia, annexitis, pelvic inflammation, bartolinitis and vaginitis.

Compounds of the invention are also useful in the treatment of ophthalmic diseases such as retinitis, retinopathies, uveitis and acute injury to the eye tissue, macular degeneration or glaucoma, conjunctivitis.

10 Compounds of the invention are also useful in the treatment of obesity.

Compounds of the invention are also useful for the treatment of all other conditions mediated by the inhibition of voltage gated sodium channels and/or voltage gated calcium channels and/or MAO-B.

It will be appreciated that the compounds of the invention may
15 advantageously be used in conjunction with one or more other therapeutic agents. Examples of suitable agents for adjunctive therapy include L-Dopa or a dopamine agonists, a 5HT_{1B/1D} agonist, such as a triptan (e.g. sumatriptan or naratriptan); an adenosine A₁ agonist; an EP ligand; an NMDA modulator, such as a glycine antagonist; a substance P antagonist (e.g. an NK1
20 antagonist); a cannabinoid; acetaminophen or phenacetin; a 5-lipoxygenase inhibitor; a leukotriene receptor antagonist; a DMARD (e.g. methotrexate); gabapentin and related compounds; a tricyclic antidepressant (e.g. amitriptyline); a neurone stabilising antiepileptic drug; a monoaminergic uptake inhibitor (e.g. venlafaxine); a matrix metalloproteinase inhibitor; a
25 nitric oxide synthase (NOS) inhibitor, such as an iNOS or an nNOS inhibitor; an inhibitor of the release, or action, of tumor necrosis factor alpha; an antibody therapy, such as monoclonal antibody therapy; an antiviral agent, such as a nucleoside inhibitor (e.g. lamivudine) or an immune system

modulator (e.g. interferon); an analgesic, such as a cyclooxygenase-2 inhibitor; a local anaesthetic; a stimulant, including caffeine; an H₂-antagonist (e.g. ranitidine); a proton pump inhibitor (e.g. omeprazole); an antacid (e.g. aluminium or magnesium hydroxide); an antifatulent (e.g. semethicone); a
5 decongestant (e.g. phenylephrine, phenylpropanolamine, pseudoephedrine, oxymetazoline, epinephrine, naphazoline, xylometazoline, propylhexedrine, or levo-desoxyephedrine, naphazoline, xylometazoline, propylhexedrine, or levo-desoxyephedrine); antitussive (e.g. codeine, hydrocodone, carmiphen, carbetapentane, or dextramethorphan); a diuretic; or a sedating or non-
10 sedating antihistamine. It is to be understood that the present invention covers the use of a compound of formula (I) or a pharmaceutically acceptable salt thereof in combination with one or more therapeutic agents.

The compounds of the present invention are useful in human and veterinary medicine. It is to be understood that reference to treatment includes
15 both treatments of established symptoms and prophylactic treatment, unless explicitly stated otherwise.

The benzylaminoalkylene heterocycles derivatives of formula I as above defined can be administered as the "active ingredient" of a pharmaceutically acceptable composition which can be prepared by
20 conventional procedures, for instance by mixing the active ingredient with pharmaceutically acceptable, therapeutically inert organic and/or inorganic carrier materials.

The composition comprising the above defined benzylaminoalkylene heterocycles derivatives can be administered in a variety forms, e.g. orally, in
25 the form of tablets, troches, capsules, sugar or film coated tablets, liquid solutions, emulsions or suspensions; rectally, in the form of suppositories; parenterally, e.g. by intramuscular or intravenous injection or infusion; and transdermally.

Suitable pharmaceutically acceptable, therapeutically inert organic and/or inorganic carrier materials useful in the preparation of such composition include, for example, water, gelatin, gum arabic, lactose, starch, cellulose, magnesium stearate, talc, vegetable oils, polyalkyleneglycols and the like. The composition comprising the benzylaminoalkylene heterocycles derivatives of formula I as above defined can be sterilized and may contain further well known components, such as, for example, preservatives, stabilizers, wetting or emulsifying agents, e.g. paraffin oil, mannide monooleate, salts to adjust osmotic pressure, buffers and the like.

For example, the solid oral forms may contain, together with the active ingredient, diluents, e.g. lactose, dextrose, saccharose, cellulose, corn starch or potato starch; lubricants, e.g. silica, talc, stearic acid, magnesium or calcium stearate, and/or polyethylene glycols; binding agents, e.g. starches, arabic gums, gelatin, methylcellulose, carboxymethylcellulose or polyvinyl pyrrolidone; disgregating agents, e.g. a starch, alginic acid, alginates or sodium starch glycolate; effervescing mixtures; dyestuffs; sweeteners; wetting agents such as lecithin, polysorbates, laurylsulphates; and, in general, non-toxic and pharmacologically inactive substances used in pharmaceutical formulations. Said pharmaceutical preparations may be manufactured in known manner, for example, by means of mixing, granulating, tableting, sugar-coating, or film-coating processes.

The oral formulations comprise sustained release formulations that can be prepared in conventional manner, for instance by applying an enteric coating to tablets and granules.

The liquid dispersion for oral administration may be e.g. syrups, emulsions and suspensions.

The syrups may contain as carrier, for example, saccharose or saccharose with glycerine and/or mannitol and/or sorbitol.

Suspensions and emulsions may contain as a carrier, for example, a natural gum, agar, sodium alginate, pectin, methylcellulose, carboxymethylcellulose, or polyvinyl alcohol. The suspensions or solutions for intramuscular injections may contain, together with the active compound, a pharmaceutically acceptable carrier, e.g. sterile water, olive oil, ethyl oleate, glycols, e.g. propylene glycol, and, if desired, a suitable amount of lidocaine hydrochloride. The solutions for intravenous injections or infusion may contain as carrier, for example, sterile water or preferably they may be in the form of sterile, aqueous, isotonic saline solutions.

The suppositories may contain, together with the active ingredient, a pharmaceutically acceptable carrier, e.g. cocoa butter, polyethylene glycol, a polyoxyethylene sorbitan fatty acid ester surfactants or lecithin.

The composition comprising the benzylaminoalkylene heterocycles derivatives of formula I as above defined is generally in the form of a dose unit containing, for example, 0.1 mg to 100 mg of active ingredient per unit dosage form.

Suitable treatment is given 1, 2 or 3 times daily, depending upon clearance rate. Accordingly, the desired dose may be presented in a single dose or as divided doses administered at appropriate intervals, for example two to four or more sub-doses per day.

The pharmaceutical compositions comprising the benzylaminoalkylene heterocycles derivatives of formula I as above defined will contain, per dosage unit, e.g., capsule, tablet, powder injection, teaspoonful, suppository and the like from about 0.1 to about 500 mg of the active ingredient most preferably from 1 to 10 mg.

Optimal therapeutically effective doses to be administered may be readily determined by those skilled in the art and will vary, basically, with the strength of the preparation, with the mode of administration and with the

advancement of the condition or disorder treated. In addition, factors associated with the particular subject being treated, including subject age, weight, diet and time of administration, will result in the need to adjust the dose to an appropriate therapeutically effective level.

5 The invention is further illustrated by the following examples.

Example 1

[4-(3-Chloro-benzyloxy)-benzyl]-(3-methyl-isoxazol-5-ylmethyl)-amine

A 1 M solution of (3-methyl-isoxazol-5-yl)-methylamine (100.9 mg, 0.9 mmol) in dry methanol was added to a 0.1 M solution of 4-(3-chloro-
10 benzyloxy)-benzaldehyde (197.4 mg, 0.8 mmol), prepared as described in Example 3, in dry methanol. 4Å molecular sieves (150 mg) were added and the reaction mixture was stirred 12 hours at room temperature under nitrogen.

A 0.5 M solution of sodium borohydride (36.6 mg, 0.97 mmol) in dry
15 methanol was added and the resulting mixture was stirred overnight at room temperature.

A 20% solution of citric acid was added (pH 5) and the crude mixture was passed through a SCX cartridge (Strong Cation Exchange resin). A mixture of the desired product and (3-methyl-isoxazol-5-yl)-methylamine were recovered by elution with 3% ammonia solution in methanol.

20 After solvent evaporation, the residue was dissolved in dichloromethane (10 ml) and treated with beads of PS-benzaldehyde (227 mg, 0.25 mmol). After resin filtration, the solvent was evaporated to dryness and 215 mg of the title compound was obtained, yield 80%.

MS (ESI Pos Spray 4 kV; Skimmer 20 V; Probe 300°C): 343.01 (MH+);
25 231.1 [M-NHR]H+

¹H-NMR (CDCl₃ + D₂O + Na₂CO₃) δ: 7.43 (s br, 1H); 7.30 (m, 2H); 7.24 (d, 2H); 6.92 (d, 2H); 5.98 (s, 1H); 5.03 (s, 2H); 3.85 (s, 2H); 3.74 (s, 2H); 2.29 (s, 3H).

Example 2**(R) [4-(3-Chloro-benzyloxy)-benzyl]-(tetrahydro-furan-2-ylmethyl)-amine**

A 1 M solution of (tetrahydro-furan-2-yl)-methylamine (91.0 mg, 0.9 mmol) in dry methanol was added to a 0.1 M solution of 4-(3-chloro-benzyloxy)-benzaldehyde (197.4 mg, 0.8 mmol), prepared as described in Example 3, in dry methanol. 4Å molecular sieves (150 mg) were added and the reaction mixture was stirred 12 hours at room temperature under nitrogen.

A 0.5 M solution of sodium borohydride (36.6 mg, 0.97 mmol) in dry methanol was added and the resulting mixture was stirred overnight at room temperature.

A 20% solution of citric acid was added (pH 5) and the crude mixture was passed through a SCX cartridge (Strong Cation Exchange resin). A mixture of the desired product and (tetrahydro-furan-2-yl)-methylamine was recovered by elution with 3% ammonia solution in methanol.

After solvent evaporation, the residue was dissolved in dichloromethane (10 ml) and treated with beads of PS-benzaldehyde (227 mg, 0.25 mmol). After resin filtration, the solvent was evaporated to dryness and 180 mg of the title compound was obtained, yield 70%.

MS (ESI Pos Spray 4 kV; Skimmer 20 V; Probe 300°C): 332.0 (MH+); 231.1 [M-NHR]H+

¹H-NMR (CDCl₃ + D₂O + Na₂CO₃) δ: 7.43 (s br, 1H); 7.33-7.21 (m, 5H); 6.92 (d, 2H); 5.03 (s, 2H); 4.08-3.96 (m, 1H); 3.86-3.68 (m, 4H); 2.77-2.61 (m, 2H); 2.02-1.80 (m, 3H); 1.62-1.49 (m, 1H).

Similarly the following compounds were prepared:

(4-Pentyloxy-benzyl)-(furan-2-ylmethyl)-amine;

(4-Heptyloxy-benzyl)-(furan-2-ylmethyl)-amine;

(R) (4-Pentyloxy-benzyl)-(tetrahydro-furan-2-ylmethyl)-amine;

- (S) (4-Pentyloxy-benzyl)-(tetrahydro-furan-2-ylmethyl)-amine;
 (R) (4-Heptyloxy-benzyl)-(tetrahydro-furan-2-ylmethyl)-amine;
 (S) (4-Heptyloxy-benzyl)-(tetrahydro-furan-2-ylmethyl)-amine;
 (R) (4-Pentyloxy-benzyl)-(tetrahydro-pyran-2-ylmethyl)-amine;
 5 (S) (4-Pentyloxy-benzyl)-(tetrahydro-pyran-2-ylmethyl)-amine;
 (R) (4-Heptyloxy-benzyl)-(tetrahydro-pyran-2-ylmethyl)-amine;
 (S) (4-Heptyloxy-benzyl)-(tetrahydro-pyran-2-ylmethyl)-amine;
 (4-Benzyloxy-benzyl)-(furan-2-ylmethyl)-amine;
 (4-Benzyloxy-benzyl)-(5-methyl-furan-2-ylmethyl)-amine;
 10 [4-(3-Fluoro-benzyloxy)-benzyl]-(furan-2-ylmethyl)-amine;
 [4-(3-Chloro-benzyloxy)-benzyl]-(furan-2-ylmethyl)-amine;
 [4-(3-Fluoro-benzyloxy)-benzyl]-(5-methyl-furan-2-ylmethyl)-amine;
 [4-(3-Chloro-benzyloxy)-benzyl]-(5-methyl-furan-2-ylmethyl)-amine;
 (R) (4-Benzyloxy-benzyl)-[1-(furan-2-yl)-1-ethyl]-amine;
 15 (S) (4-Benzyloxy-benzyl)-[1-(furan-2-yl)-1-ethyl]-amine;
 [4-(3-Fluoro-benzyloxy)-benzyl]-[1-(furan-2-yl)-1-ethyl]-amine;
 [4-(3-Chloro-benzyloxy)-benzyl]-[1-(furan-2-yl)-1-ethyl]-amine;
 (R) (4-Benzyloxy-benzyl)-(tetrahydro-furan-2-ylmethyl)-amine;
 (S) (4-Benzyloxy-benzyl)-(tetrahydro-furan-2-ylmethyl)-amine;
 20 (R) [4-(3-Fluoro-benzyloxy)-benzyl]-(tetrahydro-furan-2-ylmethyl)-amine;
 (S) [4-(3-Fluoro-benzyloxy)-benzyl]-(tetrahydro-furan-2-ylmethyl)-amine;
 (S) [4-(3-Chloro-benzyloxy)-benzyl]-(tetrahydro-furan-2-ylmethyl)-amine;
 (R) [4-(3-Fluoro-benzyloxy)-benzyl]-(tetrahydro-pyran-2-ylmethyl)-amine;
 (S) [4-(3-Fluoro-benzyloxy)-benzyl]-(tetrahydro-pyran-2-ylmethyl)-amine;
 25 (R) [4-(3-Chloro-benzyloxy)-benzyl]-(tetrahydro-pyran-2-ylmethyl)-amine;
 (S) [4-(3-Chloro-benzyloxy)-benzyl]-(tetrahydro-pyran-2-ylmethyl)-amine;
 (4-Benzyloxy-benzyl)-(1,4-dioxan-2-ylmethyl)-amine;
 [4-(3-Chloro-benzyloxy)-benzyl]-(pyrido-2-ylmethyl)-amine;

- [4-(3-Chloro-benzyloxy)-benzyl]-(pyrido-3-ylmethyl)-amine;
[4-(3-Chloro-benzyloxy)-benzyl]-(pyrido-4-ylmethyl)-amine;
[4-(3-Chloro-benzyloxy)-benzyl]-(imidazol-2-ylmethyl)-amine;
[4-(3-Chloro-benzyloxy)-benzyl]-(1-methyl-imidazol-2-ylmethyl)-amine;
5 [4-(3-Chloro-benzyloxy)-benzyl]-(5-methyl-1H-triazol-2-ylmethyl)-amine;
[4-(3-Chloro-benzyloxy)-benzyl]-(4-methyl-thiazol-2-ylmethyl)-amine;
(4-Benzyloxy-benzyl)-(isoxazol-5-ylmethyl)-amine;
[4-(3-Fluoro-benzyloxy)-benzyl]-(isoxazol-5-ylmethyl)-amine;
[4-(3-Chloro-benzyloxy)-benzyl]-(isoxazol-5-ylmethyl)-amine;
10 (4-Benzyloxy-benzyl)-(3-methyl-isoxazol-5-ylmethyl)-amine;
[4-(3-Fluoro-benzyloxy)-benzyl]-(3-methyl-isoxazol-5-ylmethyl)-amine;
(4-Benzyloxy-benzyl)-(oxazol-2-ylmethyl)-amine;
[4-(3-Fluoro-benzyloxy)-benzyl]-(oxazol-2-ylmethyl)-amine;
[4-(3-Chloro-benzyloxy)-benzyl]-(oxazol-2-ylmethyl)-amine;
15 (4-Benzyloxy-benzyl)-(oxazol-5-ylmethyl)-amine;
[4-(3-Fluoro-benzyloxy)-benzyl]-(oxazol-5-ylmethyl)-amine;
[4-(3-Chloro-benzyloxy)-benzyl]-(oxazol-5-ylmethyl)-amine;
(4-Benzyloxy-benzyl)-(thiophen-2-ylmethyl)-amine;
[2-(3-Chloro-benzyloxy)-benzyl]-(thiophen-2-ylmethyl)-amine;
20 [2-(3-Fluoro-benzyloxy)-benzyl]-(thiophen-2-ylmethyl)-amine;
[3-(3-Chloro-benzyloxy)-benzyl]-(thiophen-2-ylmethyl)-amine;
[3-(3-Fluoro-benzyloxy)-benzyl]-(thiophen-2-ylmethyl)-amine;
[4-(3-Chloro-benzyloxy)-benzyl]-(thiophen-2-ylmethyl)-amine;
[4-(3-Fluoro-benzyloxy)-benzyl]-(thiophen-2-ylmethyl)-amine;
25 [2-(3-Fluoro-benzyloxy)-benzyl]-(benzo[b]furan-2-ylmethyl)-amine;
[3-(3-Fluoro-benzyloxy)-benzyl]-(benzo[b]furan-2-ylmethyl)-amine;
[4-(3-Fluoro-benzyloxy)-benzyl]-(benzo[b]furan-2-ylmethyl)-amine;
(R) [4-(3-Fluoro-benzyloxy)-benzyl]-(dihydro-benzo[b]furan-2-ylmethyl)-

amine;

(S) [4-(3-Fluoro-benzyloxy)-benzyl]-(dihydro-benzo[b]furan-2-ylmethyl)-

amine;

[4-(3-Chloro-benzyloxy)-benzyl]-(benzimidazol-2-ylmethyl)-amine;

5 **Example 3**

4-(3-Chloro-benzyloxy)-benzaldehyde

A 0.5 M solution of 1-bromomethyl-3-chloro-benzene (102.7 mg, 0.5 mmol) in acetone was added to a suspension of 4-hydroxy-benzaldehyde (67.2 mg, 0.55 mmol), K_2CO_3 (207.3 mg, 1.5 mmol) and KI (10.8 mg, 0.065 mmol) 10 in acetone (7 ml). The reaction mixture was stirred for 2 hours at 70°C.

After cooling, the solid residue was filtered off and the solvent was evaporated under vacuum. The residue was dissolved in ethyl acetate and the organic layer washed twice with 1 M NaOH and dried over Na_2SO_4 . The title compound was obtained (85 mg) after evaporating the solvent under vacuum, 15 yielding 96.6%.

Example 4

***In vitro* MAO-A and MAO-B enzyme activities assay**

Membrane preparations (crude mitochondrial fraction): male Wistar rats (Harlan, Italy – 175-200 g) were sacrificed under light anaesthesia and 20 brains were rapidly removed and homogenized in 8 vol. of ice-cold 0.32 M sucrose buffer containing 0.1 M EDTA, pH 7.4. The crude homogenate was centrifuged at 2220 rpm for 10 min and the supernatant recovered. The pellet was homogenized and centrifuged again and the two supernatants were pooled and centrifuged at 9250 rpm for 10 min, +4°C. The pellet was resuspended in 25 fresh buffer and centrifuged at 11250 rpm for 10 min, +4°C. The resulting pellet was stored at -80°C until use.

***In vitro* enzyme activities assay:** the enzyme activities were assessed with a radioenzymatic assay using the selective substrates ^{14}C -serotonin (5-HT)

29

and ^{14}C -phenylethylamine (PEA) for MAO-A and MAO-B, respectively.

The mitochondrial pellet (500 μg protein) was resuspended in 0.1 M phosphate buffer pH 7.4 and 500 μl was added to 50 μl of the test compound or buffer for 30 min at 37°C (preincubation) then the substrate (50 μl) was
5 added. The incubation was carried out for 30 min at 37°C (^{14}C -5-HT, 5 μM) or for 10 min at 37°C (^{14}C -PEA, 0.5 μM).

The reaction was stopped by adding 0.2 ml HCl or perchloric acid. After centrifugation, the deaminated metabolites were extracted with 3 ml of diethylether (5-HT) or toluene (PEA) and the radioactive organic phase
10 measured by liquid scintillation spectrometry at 90% efficiency. Radioactivity in the eluate indicates the production of neutral and acidic metabolites formed as a result of MAO activity.

The enzymatic activity was expressed as nmoles of substrate transformed/mg protein/min. The activity of MAO in the sample was
15 expressed as a percentage of control activity in the absence of inhibitors after subtraction of appropriate blank values.

The drug inhibition curves were obtained from at least eight different concentrations, each in duplicate (10^{-10} to 10^{-5} M) and the IC_{50} values (the drug concentration inhibiting 50% enzyme activity) with confidence intervals
20 determined using non linear regression analysis (best fitting aided-computer program).

Example 5

In vitro MAO-B inhibition reversibility studies

To investigate whether the test compound was an irreversible or a
25 reversible MAO-B inhibitor, the inhibition of the enzymatic activity was assessed using two experimental protocols:

- a) *time-dependent experiments:*

the time-dependent association kinetics were measured as the IC_{50}

30

values after 0 or 30 min enzyme-inhibitor incubation. For mechanism-based irreversible inhibitors that act by blocking the enzyme catalytic site, the inhibitory potency increases with incubation time. The absence of significant difference between IC_{50} with or without preincubation is indicative for reversible inhibitors.

- *b) washing experiments:*

rat brain mitochondria (enzyme) and test compound (inhibitor) were preincubated for 30 min at 37°C before being subjected to repeated centrifugations and resuspensions. MAO-B activity was assessed immediately and after each centrifugation resuspension cycles (1-5). Percentage inhibition was calculated with respect to samples treated in the same manner but in the absence of inhibitor.

The recovery of MAO-B activity after washings of the enzyme-test compound complex is indicative for reversible inhibitors.

Example 6

***Ex vivo* MAO-B inhibition**

Test compounds were administered orally to male C57BL mice (Harlan, Italy, 25-27 g) at the single dose of 20 mg/Kg. At different time intervals (1, 2, 4, 8 and 24 h), animals were sacrificed, brains removed, cortices dissected out and stored at -80°C. Crude homogenates (0.5%) were prepared in 0.1 M phosphate buffer pH 7.4 and freshly used. MAO-A and MAO-B activity were assessed as above described.

Example 7

***In vitro* binding of 3H -batrachotoxin in rat brain membranes**

Membrane preparations (P2 fraction) Male Wistar rats (Harlan, Italy - 175-200 g) were sacrificed under light anaesthesia and brains were rapidly removed and cortex was dissected out, homogenized in 10 vol. of ice-cold 0.25 M sucrose buffer (50 mM Tris·HCl, pH 7.4). The crude homogenate was

centrifuged at 3250 rpm for 10 min and the supernatant recovered. The pellet was homogenized and centrifuged again and the two supernatants were pooled and centrifuged at 14750 rpm for 10 min, +4°C. The resulting pellet was stored at -20°C until use.

5 Binding assay. The pellet was resuspended in 50 mM Hepes buffer, pH 7.4 containing 0.8 mM MgSO₄, 5.4 mM KCl, 5.5 mM glucose and 130 mM choline using Polytron PT10. The binding assay was carried out in 0.25 ml final volume containing 50 µl of membrane preparation (ca 200 µg of protein), 50 µl of ³H-batrachotoxin ligand (10 nM), 50 µl of TTX (1 µM), 50 µl of
10 scorpion toxin (37.5 µg/ml) and 50 µl of test compound or buffer or 300 µM of veratridine to determine non specific binding. The binding assay was performed at 37°C for 30 min and stopped by rapid filtration under vacuum through Whatman GF/B glass fiber filters. Filters (pre-soaked with polyethylenimine 0.1%) were washed with 3x5 ml of ice-cold buffer and
15 placed in picovials containing scintillation cocktail (Filter Count, Packard). Radioactivity bound was measured by liquid scintillation spectrometry at 45% efficiency.

Data analysis. The IC₅₀ was calculated from displacement curves by computer program LIGAND (McPherson, J. Pharmacol. Methods, 14, 213.
20 1985). The displacement curves were obtained using at least 9 concentrations, each in duplicate, covering a 100000 fold range.

Example 8

In vitro binding of ³H-nitrendipine in rat brain membranes

Membrane preparations Male Wistar rats (Harlan, Italy - 175-200 g)
25 were sacrificed under light anaesthesia and brains were rapidly removed and cortex was dissected out, homogenized in 10 vol. of ice-cold 50 mM Tris•HCl, pH 7.7 using Polytron PT10. The crude homogenate was centrifuged at 50000xg for 10 min. The pellet was homogenized and centrifuged twice in

32

fresh buffer at 50000xg for 10 min, +4°C. The resulting pellet was stored at -20°C until use.

Binding assay. The pellet was resuspended in 50 mM Tris-HCl, pH 7.7 using Polytron PT10. The binding assay was carried out in 1 ml final volume containing 900 µl of membrane preparation (ca 700 µg of protein), 50 µl of ³H-nitrendipine (0.15 nM), and 50 µl of test compound or buffer or 1 µM of nifedipine to determine non specific binding. The binding assay was performed at 25°C for 45 min and stopped by rapid filtration under vacuum through Whatman GF/B glass fiber filters. Filters were washed with 3x5 ml of ice-cold buffer and placed in picovials containing scintillation cocktail (Filter Count, Packard). Radioactivity bound was measured by liquid scintillation spectrometry at 45% efficiency.

Data analysis. The IC₅₀ was calculated from displacement curves by computer program LIGAND (McPherson, J. Pharmacol. Methods, 14, 213. 1985). The displacement curves were obtained using at least 9 concentrations, each in duplicate, covering a 100000 fold range.

Example 9

Calcium influx assay

IMR32 human neuroblastoma cells constitutively possess both L and N type channels. Under differentiating conditions, IMR32 preferentially express on the membrane surface N- type calcium channels. The remaining L-type calcium channels were blocked using the selective L type blocker, nifedipine. In these experimental conditions, only N type channels can be detected.

IMR32 cells were differentiated using 1 mM dibutyryl-cAMP and 2.5 µM bromodeoxyuridine for 8 days (4 times) in 225 cm² flask, then detached, seeded at 200,000 cells/well on 96 polyllysine-coated plates and further incubated for 18-24 h in the presence of differentiating buffer before use.

The Ca²⁺ Kit Assay (Molecular Devices), based on a fluorescent

33

calcium indicator 485-535 nm wavelength, was used.

Differentiated cells were incubated with dye loading for 30 min at 37°C then, nifedipine alone (1 μ M) or in the presence of ω -conotoxin or test compounds were added for further 15 min.

- 5 The fluorescence (485-535 nm) was measured before and after (30-40 sec) the automated injection of 100 mM KCl depolarizing solution using a Victor plate reader (Perkin Elmer).

Example 10

Maximal electroshock test in rats and mice

- 10 Wistar rats received an electroshock (160 mA for 0.2 s with a pulse train of 60 Hz having a pulse duration of 0.4 ms; ECT unit model 7801, Ugo Basile, Comerio, Italy) through intra-aural clip electrodes sufficient to produce a hindlimb tonic extensor response in at least 97% of control animals.

- 15 Mice received a 28 mA shock of 0.7 s with a pulse train of 80 Hz having a pulse duration of 0.4 ms. Several doses of the test compound and standard AEDs were administered to groups of 10 to 20 mice or rats per dose in a volume of 5 ml/kg 60 min p.o. or 30 min i.p. before induction of MES to calculate ED₅₀ values. The complete suppression of the hindlimb tonic extensor component of seizures was taken as evidence of anticonvulsant
20 activity.

Example 11

Mice Formalin Test

- 25 According to a modified protocol from Rosland et al., (1990) mice were injected subcutaneously (s.c.) with 20 μ l of 2.7% solution of formalin into the plantar surface of left hindpaw and placed immediately into clear PVC observation chambers (23 x 12 x 13 cm). Pain behaviour was quantified by counting the cumulative licking time (s) of the injected paw. Measurements were taken during the early phase (0-5 min) and late phase (30-40 min) after

34

formalin injection (Tjolsen et al. 1992).

The test compound was administered p.o. 15 min before formalin injection in a volume of 10 ml/kg body weight to groups of 10 mice per dose. Control group was treated with vehicle.

5 **Example 12**

Carragenan model of inflammation

Male Wistar rats of 175-200 grams were used.


10 The left hind paw was injected with 100 μ l of carrageenan (2% w/v in saline). Compounds of the invention (30 mg/kg), indomethacin (5 mg/kg) or control vehicle (such as distilled water) were orally administered 1 h before carrageenan injection. The paw volume was measured with a plethysmometer (Ugo Basile) immediately before (basal) and 1, 2, 3, 4 and 5 h after the carrageenan injection.

15

35

CLAIMS

1. Compounds of general formula I:



I

wherein

X is oxygen or sulphur or a NR^7 group;

R^1 is $\text{C}_3\text{-C}_8$ alkyl, or $\text{C}_1\text{-C}_8$ alkyl substituted by phenoxy or phenyl, both phenoxy or phenyl being optionally substituted by one or more fluoro, chloro, trifluoromethyl, $\text{C}_1\text{-C}_6$ alkyl, hydroxyl, $\text{C}_1\text{-C}_6$ alkoxy;

R^2, R^3 are independently hydrogen, $\text{C}_1\text{-C}_6$ alkyl, halogen, trifluoromethyl, hydroxy or $\text{C}_1\text{-C}_6$ alkoxy;

R^4 is hydrogen, $\text{C}_1\text{-C}_8$ alkyl;

R^5, R^6 are independently hydrogen, $\text{C}_1\text{-C}_3$ alkyl, optionally substituted by hydroxy or phenyl;

R^7 is hydrogen or straight or branched $\text{C}_1\text{-C}_3$ alkyl;


Het is a five to seven membered, saturated or unsaturated heteromonocyclic or an eight to ten membered, saturated or unsaturated heterobicyclic group, containing one or more heteroatoms chosen independently from nitrogen, oxygen and sulphur, said mono- or bicyclic groups being optionally substituted by $\text{C}_1\text{-C}_6$ alkyl, halogen, hydroxyl or $\text{C}_1\text{-C}_6$ alkoxy;

and the pharmaceutically acceptable salts thereof, with the proviso that:

36

- Het cannot be indole, benzo[b]furan, benzo[b]thiophen, chroman, when R^5 and R^6 are both hydrogen, or 2-pyridyl;
- when R^1 is unsubstituted or substituted benzyl, and R^2 and R^3 are hydrogen, halogen or alkoxy, R^4 is other than hydrogen;
- 5 • when R^1 is propyl or butyl, R^2 , R^3 , R^5 and R^6 are hydrogen and R^4 is hydrogen, methyl or ethyl, Het cannot be 1,4-benzodioxan;
- when X- R^1 is a para butyloxy group and R^2 , R^3 , R^4 , R^5 and R^6 are hydrogen, Het cannot be 2-thiophenyl, or 4-(2,2'-dimethyl)-pyranyl;
- 10 • when X- R^1 is an ortho heptyloxy or octyloxy group and R^2 , R^3 , R^4 , R^5 and R^6 are hydrogen, Het cannot be 2-furyl;
- when X- R^1 is an ortho O-(CH₂)_m-p-CF₃-phenyl group, where m is an integer from 1 to 3, and R^2 , R^3 , R^5 and R^6 are hydrogen, Het cannot be pyridyl.

15 2. The use of the compounds of formula I



wherein

- 20 R^1 is C₃-C₈ alkyl, or C₁-C₃ alkyl substituted by phenoxy or phenyl, both phenoxy or phenyl being optionally substituted by one or more fluoro, chloro, trifluoromethyl, C₁-C₆ alkyl, hydroxyl, C₁-C₆ alkoxy;
- R^2 , R^3 are independently hydrogen, C₁-C₆ alkyl, halogen, trifluoromethyl, hydroxy or C₁-C₆ alkoxy;
- 25 R^4 is hydrogen, C₁-C₈ alkyl;

R^5, R^6 are independently hydrogen, C_1-C_3 alkyl, optionally substituted by hydroxy or phenyl;

R^7 is hydrogen or straight or branched C_1-C_3 alkyl;

X is oxygen or sulphur or a NR^7 group;

5 Het is a five to seven membered, saturated or unsaturated heteromonocyclic or an eight to ten membered, saturated or unsaturated heterobicyclic group, containing one or more heteroatoms chosen independently from nitrogen, oxygen and sulphur, said mono- or bicyclic groups being optionally substituted by C_1-C_6 alkyl, halogen, hydroxyl or C_1-C_6 alkoxy;

10 and the pharmaceutically acceptable salts thereof, with the proviso that:

- Het cannot be an indole, chroman when R^5 and R^6 are both hydrogen;
- when X- R^1 is an ortho- $O-(CH_2)_m$ -p- CF_3 -phenyl group, where m is an integer from 1 to 3, and R^2, R^3, R^5 and R^6 are hydrogen, Het cannot be pyridyl

and the pharmaceutically acceptable salts or prodrug thereof, for the preparation of a medicament having sodium and/or calcium channel modulating activity and/or selective MAO-B inhibiting activity and therefore useful in preventing, alleviating and curing a wide range of pathologies, including, but not limited to, neurological, psychiatric, cardiovascular, inflammatory, ophthalmic, urologic, metabolic and gastrointestinal diseases

20 3. The use of compounds according to claim 2 wherein R^1 is benzyl or C_5-C_8 alkyl, R^4, R^5 and R^6 are hydrogen or C_1-C_3 alkyl, X is oxygen and Het is furan, tetrahydrofuran, isoxazol, oxazol, thiophen, pyran, dioxane, unsubstituted or substituted by C_1-C_3 alkyl.

4. The use of a compound according to claim 2 selected from the group consisting of:

38

- (4-Pentyloxy-benzyl)-(furan-2-ylmethyl)-amine;
(4-Heptyloxy-benzyl)-(furan-2-ylmethyl)-amine;
(R) (4-Pentyloxy-benzyl)-(tetrahydro-furan-2-ylmethyl)-amine;
(S) (4-Pentyloxy-benzyl)-(tetrahydro-furan-2-ylmethyl)-amine;
5 (R) (4-Heptyloxy-benzyl)-(tetrahydro-furan-2-ylmethyl)-amine;
(S) (4-Heptyloxy-benzyl)-(tetrahydro-furan-2-ylmethyl)-amine;
(R) (4-Pentyloxy-benzyl)-(tetrahydro-pyran-2-ylmethyl)-amine;
(S) (4-Pentyloxy-benzyl)-(tetrahydro-pyran-2-ylmethyl)-amine;
(R) (4-Heptyloxy-benzyl)-(tetrahydro-pyran-2-ylmethyl)-amine;
10 (S) (4-Heptyloxy-benzyl)-(tetrahydro-pyran-2-ylmethyl)-amine;
(4-Benzyloxy-benzyl)-(furan-2-ylmethyl)-amine;
(4-Benzyloxy-benzyl)-(5-methyl-furan-2-ylmethyl)-amine;
[4-(3-Fluoro-benzyloxy)-benzyl]-(furan-2-ylmethyl)-amine;
[4-(3-Chloro-benzyloxy)-benzyl]-(furan-2-ylmethyl)-amine;
15 [4-(3-Fluoro-benzyloxy)-benzyl]-(5-methyl-furan-2-ylmethyl)-amine;
[4-(3-Chloro-benzyloxy)-benzyl]-(5-methyl-furan-2-ylmethyl)-amine;
(R) (4-Benzyloxy-benzyl)-[1-(furan-2-yl)-1-ethyl]-amine;
(S) (4-Benzyloxy-benzyl)-[1-(furan-2-yl)-1-ethyl]-amine;
[4-(3-Fluoro-benzyloxy)-benzyl]-[1-(furan-2-yl)-1-ethyl]-amine;
20 [4-(3-Chloro-benzyloxy)-benzyl]-[1-(furan-2-yl)-1-ethyl]-amine;
(R) (4-Benzyloxy-benzyl)-(tetrahydro-furan-2-ylmethyl)-amine;
(S) (4-Benzyloxy-benzyl)-(tetrahydro-furan-2-ylmethyl)-amine;
(R) [4-(3-Fluoro-benzyloxy)-benzyl]-(tetrahydro-furan-2-ylmethyl)-amine;
(S) [4-(3-Fluoro-benzyloxy)-benzyl]-(tetrahydro-furan-2-ylmethyl)-amine;
25 (R) [4-(3-Chloro-benzyloxy)-benzyl]-(tetrahydro-furan-2-ylmethyl)-amine;
(S) [4-(3-Chloro-benzyloxy)-benzyl]-(tetrahydro-furan-2-ylmethyl)-amine;
(R) [4-(3-Fluoro-benzyloxy)-benzyl]-(tetrahydro-pyran-2-ylmethyl)-amine;
(S) [4-(3-Fluoro-benzyloxy)-benzyl]-(tetrahydro-pyran-2-ylmethyl)-amine;

- (R) [4-(3-Chloro-benzyloxy)-benzyl]-(tetrahydro-pyran-2-ylmethyl)-amine;
 (S) [4-(3-Chloro-benzyloxy)-benzyl]-(tetrahydro-pyran-2-ylmethyl)-amine;
 (4-Benzyloxy-benzyl)-(1,4-dioxan-2-ylmethyl)-amine;
 [4-(3-Chloro-benzyloxy)-benzyl]-(pyrido-2-ylmethyl)-amine;
 5 [4-(3-Chloro-benzyloxy)-benzyl]-(pyrido-3-ylmethyl)-amine;
 [4-(3-Chloro-benzyloxy)-benzyl]-(pyrido-4-ylmethyl)-amine;
 [4-(3-Chloro-benzyloxy)-benzyl]-(imidazol-2-ylmethyl)-amine;
 [4-(3-Chloro-benzyloxy)-benzyl]-(1-methyl-imidazol-2-ylmethyl)-amine;
 [4-(3-Chloro-benzyloxy)-benzyl]-(5-methyl-1H-triazol-2-ylmethyl)-amine;
 10 [4-(3-Chloro-benzyloxy)-benzyl]-(4-methyl-thiazol-2-ylmethyl)-amine;
 (4-Benzyloxy-benzyl)-(isoxazol-5-ylmethyl)-amine;
 [4-(3-Fluoro-benzyloxy)-benzyl]-(isoxazol-5-ylmethyl)-amine;
 [4-(3-Chloro-benzyloxy)-benzyl]-(isoxazol-5-ylmethyl)-amine;
 (4-Benzyloxy-benzyl)-(3-methyl-isoxazol-5-ylmethyl)-amine;
 15 [4-(3-Fluoro-benzyloxy)-benzyl]-(3-methyl-isoxazol-5-ylmethyl)-amine;
 [4-(3-Chloro-benzyloxy)-benzyl]-(3-methyl-isoxazol-5-ylmethyl)-amine;
 (4-Benzyloxy-benzyl)-(oxazol-2-ylmethyl)-amine;
 [4-(3-Fluoro-benzyloxy)-benzyl]-(oxazol-2-ylmethyl)-amine;
 [4-(3-Chloro-benzyloxy)-benzyl]-(oxazol-2-ylmethyl)-amine;
 20 (4-Benzyloxy-benzyl)-(oxazol-5-ylmethyl)-amine;
 [4-(3-Fluoro-benzyloxy)-benzyl]-(oxazol-5-ylmethyl)-amine;
 [4-(3-Chloro-benzyloxy)-benzyl]-(oxazol-5-ylmethyl)-amine;
 (4-Benzyloxy-benzyl)-(thiophen-2-ylmethyl)-amine;
 [2-(3-Chloro-benzyloxy)-benzyl]-(thiophen-2-ylmethyl)-amine;
 25 [2-(3-Fluoro-benzyloxy)-benzyl]-(thiophen-2-ylmethyl)-amine;
 [3-(3-Chloro-benzyloxy)-benzyl]-(thiophen-2-ylmethyl)-amine;
 [3-(3-Fluoro-benzyloxy)-benzyl]-(thiophen-2-ylmethyl)-amine;
 [4-(3-Chloro-benzyloxy)-benzyl]-(thiophen-2-ylmethyl)-amine;

40


- [4-(3-Fluoro-benzyloxy)-benzyl]-(thiophen-2-ylmethyl)-amine;
 [2-(3-Fluoro-benzyloxy)-benzyl]-(benzo[b]furan-2-ylmethyl)-amine;
 [3-(3-Fluoro-benzyloxy)-benzyl]-(benzo[b]furan-2-ylmethyl)-amine;
 [4-(3-Fluoro-benzyloxy)-benzyl]-(benzo[b]furan-2-ylmethyl)-amine;
 5 (R) [4-(3-Fluoro-benzyloxy)-benzyl]-(dihydro-benzo[b]furan-2-ylmethyl)-
 amine;
 (S) [4-(3-Fluoro-benzyloxy)-benzyl]-(dihydro-benzo[b]furan-2-ylmethyl)-
 amine;

- [4-(3-Chloro-benzyloxy)-benzyl]-(benzimidazol-2-ylmethyl)-amine;
 10 either as a single isomer or as a mixture of isomers thereof, and the
 pharmaceutically acceptable salts thereof.

5. A process for the preparation of a compound of formula I, as defined in
 claim 1, or a pharmaceutically acceptable salt thereof, the process comprising:

- a) a reaction of a compounds of formula II


15



II

wherein R^1 , R^2 , R^3 and X are as defined in claim 1
 with compounds of formula III, in the presence of a reducing agent

20




III

wherein R^4 , R^5 , R^6 and Het are as defined in claim 1, in the presence of
 a reducing agent or

41

- b) a reaction of a compound of formula III with a compound of formula IV,



IV

5

wherein R^1 , R^2 , R^3 and X are as defined above and Y is a halogen atom or a O-EWG group, where the EWG means an electron withdrawing group, able to transform the oxygen which they are linked to, in a good leaving group and, if desired, converting a compound of the invention into another compound of the invention and/or, if desired, converting a compound of the invention into a pharmaceutically acceptable salt and/or, if desired, converting a salt into a free compound.


6. A pharmaceutical composition containing, as an active principle, a compound of formula I, as defined in claim 1, or a pharmaceutically acceptable salt thereof in addition to a suitable carrier and/or diluent and optionally to other therapeutic agents.

7. A compound of formula I, as defined in claim 1, or a pharmaceutically acceptable salt thereof, for use as an active therapeutic substance.

42

ABSTRACT**SUBSTITUTED BENZYLAMINOALKYLENE HETEROCYCLES**

5 This invention is related to compounds of general formula I



wherein

- 10 X is oxygen or sulphur or a NR^7 group;
 R^1 is $\text{C}_3\text{-C}_8$ alkyl, or $\text{C}_1\text{-C}_8$ alkyl substituted by phenoxy or phenyl, both phenoxy or phenyl being optionally substituted by one or more fluoro, chloro, trifluoromethyl, $\text{C}_1\text{-C}_6$ alkyl, hydroxyl, $\text{C}_1\text{-C}_6$ alkoxy;
- 15 R^2, R^3 are independently hydrogen, $\text{C}_1\text{-C}_6$ alkyl, halogen, trifluoromethyl, hydroxy or $\text{C}_1\text{-C}_6$ alkoxy;
 R^4 is hydrogen, $\text{C}_1\text{-C}_8$ alkyl;
 R^5, R^6 are independently hydrogen, $\text{C}_1\text{-C}_3$ alkyl, optionally substituted by hydroxy or phenyl;
- 20 R^7 is hydrogen or straight or branched $\text{C}_1\text{-C}_3$ alkyl;
Het is a five to seven membered, saturated or unsaturated heteromonocyclic or an eight to ten membered, saturated or unsaturated heterobicyclic group, containing one or more heteroatoms chosen independently from nitrogen, oxygen and sulphur, said mono or bicyclic groups being optionally
- 25 substituted by $\text{C}_1\text{-C}_6$ alkyl, halogen, hydroxyl or $\text{C}_1\text{-C}_6$ alkoxy;

43

and the pharmaceutically acceptable salts or prodrug thereof, that are active as sodium and/or calcium channel modulators and/or as selective MAO-B inhibitors and therefore useful in preventing, alleviating and curing a wide range of pathologies, including, but not limited to, neurological, psychiatric, cardiovascular, inflammatory, ophthalmic, urologic, metabolic and gastrointestinal diseases, where the above mechanisms have been described as playing a pathological role.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.